

Technical University of Denmark



Multi-omic profiling of EPO-producing CHO cell panel reveals metabolic adaptation to heterologous protein production

Ley, Daniel; Kazemi Seresht, Ali; Engmark, Mikael; Magdenoska, Olivera; Nielsen, Kristian Fog; Kildegaard, Helene Fastrup; Andersen, Mikael Rørdam

Publication date:
2016

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Ley, D., Kazemi Seresht, A., Engmark, M., Magdenoska, O., Nielsen, K. F., Kildegaard, H. F., & Andersen, M. R. (2016). Multi-omic profiling of EPO-producing CHO cell panel reveals metabolic adaptation to heterologous protein production. Abstract from Cell Culture Engineering XV, La Quinta, CA, United States.

DTU Library
Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

MULTI-OMIC PROFILING OF EPO PRODUCING CHO CELL PANEL REVEALS METABOLIC ADAPTATION TO HETEROLOGOUS PROTEIN PRODUCTION

Daniel Ley, Technical University of Denmark, Department of Systems Biology
Daley@bio.dtu.dk

Ali Kazemi Seresht, Novo Nordisk A/S

Mikael Engmark, Technical University of Denmark, Department of Systems Biology

Olivera Magdenoska, Technical University of Denmark, Department of Systems Biology

Kristian Fog Nielsen, Technical University of Denmark, Department of Systems Biology

Helene Fastrup Kildegaard, Novo Nordisk Foundation Center for Biosustainability

Mikael Rørdam Andersen, Technical University of Denmark, Department of Systems Biology

Key Words: Chinese hamster ovary, metabolic adaptation, chemostat, metabolomics, transcriptomics

The Chinese hamster ovary (CHO) cell line is the predominant mammalian cell factory for production of therapeutic glycoproteins. In this work, we aimed to study bottlenecks in the secretory pathway associated with the production of human erythropoietin (EPO) in CHO cells. In connection to this, we discovered indications of metabolic adaptation of the amino acid catabolism in favor of heterologous protein production. We established a panel of stably EPO expressing CHO-K1 clones spanning a 25-fold productivity range and characterized the clones in batch and chemostat cultures. For this, we employed a multi-omic physiological characterization including metabolic footprinting of amino acids, metabolite fingerprinting of glycolytic intermediates, NAD(P)H-/NAD(P)⁺ and adenosine nucleotide phosphates. We used qPCR, qRT-PCR, western blots and Affymetrix CHO microarrays to assess EPO gene copy numbers, EPO gene expression, intracellular protein levels and genome-wide differential gene expression analysis of genes functionally related to secretory protein processing, respectively. Finally, we generated a network reconstruction of the amino acid catabolism in CHO cells. The reconstruction was utilized as a platform for interpretation of differential gene expression data in a biological meaningful manner. To identify bottlenecks in the protein secretory pathway, we compared EPO gene copy numbers, EPO gene expression levels, intracellular EPO retention and extracellular EPO levels for a high and low producing clone during chemostat culture. The EPO productivity levels were not reflected in EPO gene load, EPO gene expression or intracellular protein retention, indicating that these processes were not limiting EPO productivity. The global gene expression analysis did not identify significant differentially expressed genes related to secretory protein processing. However, when inspecting the gene expression landscape of the amino acid catabolism, we observed an apparent adaptation in favor of EPO production. That is, we discovered that the gene expression levels of amino acid catabolic genes had adapted to preserve the most abundant amino acids in EPO in the high producing clone relative to the low producing clone. Based on these data, we speculate that the amino acid metabolism in CHO cells may undergo adaptation in favor of heterologous protein production during long-term cultivation.